Bulletin of the British Myriapod Group 6

LITHOBIUS TENEBROSUS MEINERT FROM ABERYSTWYTH, CARDIGANSHIRE

A N Keay 46 Albany Road, Newport, Isle of Wight PO30 5JA

<u>Lithobius tenebrosus</u> was first recorded from Britain by R S Bagnall in 1913 from County Durham and again from Cornwall in 1945 by F A Turk (both as <u>L. nigrifons</u>). Neither of these early records have been confirmed and Dr Eason has some doubt of the validity of the Bagnall record (pers.comm.,1988).

The April 1988 field trip of the BMG/BISG to Constitution Hill, Aberystwyth (22/58-81-) has provided the first confirmed record of <u>L.tenebrosus</u> from Britain. The specimen was found in a crevice in a sea cliff at about 10 metres. Other species present at the site were <u>Strigamia crassipes</u>, <u>Cryptops hortensis</u>, <u>Lithobius forficatus</u> and <u>Lithobius melanops</u>.

Description

A pseudomaturus female, light chestnut brown (slightly mottled)

Length: 9.5mm Breadth: 1.0mm Antennae: 35 articles (Fig.1)

Head: Marginal ridge interrupted and slightly projecting laterally (Fig.1)

Ocelli (1+13) are more numerous than would be expected in a pseudomaturus of other species of a comparable size.

Forcipular segment with 2+2 prosternal teeth in almost a straight line. Prosternal teeth well developed and the internal pair slightly larger than the external pair. Parodontal spine slender (Fig. 3).

Trunk: T9, T11 and T13 with small prosterior projections (the projections on T13 are almost as small as those on T9 and not larger as in <u>L.melanops</u> etc.)(Fig.2).

Female genitalia with 2+2 gonopod spurs (Fig.4) and a narrow claw.

Coxal pores: Round with 3 on each leg. (3-6 in mature specimens)

Spinulation (see Figs 5 and 6)

DaC absent

15 DaP absent

15 VaF present

15 VmC absent

15th legs without accessory claw.


<u>Lithobius tenebrosus</u> is known from Austria, Germany, Scandinavia and Switzerland.


References

Eason, E.H. (1964). Centipedes of the British Isles. London, Warne. Eason, E.H. (1981). A review of the north-west European species of Lithobiomorpha with a revised key to their identification. Zool. J. Linn. Soc. (1982) 74:9-33.

Acknowledgements

My grateful thanks to Ted Eason and Tony Barber for their prompt confimation of the specimen and to Dick Jones for his wonderful artwork.





Figs 1-6 <u>Lithobius tenebrosus</u> pseudomaturus female from Aberystwyth:

- 1. Head and first tergite, dorsal
- 2. Tergites 8-15
- 3. Forcipular prosternal teeth, ventral
- 4. Gonopods ventral
- 5. 15th leg, ventral
- 6. 15th leg, dorsal